



Leptastrea purpurea

Crust coral

Montano, S.

Citation: Montano, S., 2022. *Leptastrea purpurea*, Crust coral. The Maldives Red List of Threatened Species.

A. Background Information

1. Assessment Information:

Assessor Name	Simone Montano
Date of Assessment	17/02/2022
Contributors	Ahmed Basheer, Hana Amir, Phillipa Roe, Ilham Mohamed
Institutional contributors	Ministry of Environment, Climate Change and Technology; Maldives Marine Research Institute; IUCN Project REGENERATE; Maldives Allen Coral Atlas; Noo Raajje; Six Senses Laamu, MaRHE center
Facilitators	James Tallant
Reviewers	Janice Chanson, Maldives Marine Research Institute


2. Taxonomic Information:

Scientific Name	<i>Leptastrea purpurea</i> (Dana, 1846)	
Common Name (English)	Crust coral	
Common Name (Dhivehi)		
Taxonomy	Order	Scleractinia
	Sub-order	-
	Family	Leptastreaeidae
Taxonomic Notes	-	

3. Geographic Range:

Summary of Global Distribution	In the Indo-West Pacific, this species is found in the Red Sea and Gulf of Aden, southwest Indian Ocean, NW Indian Ocean and Arabian/Iranian Gulf, Northern Indian Ocean, Central Indo-Pacific, Australia, Southeast Asia, Japan and East China Sea, oceanic West Pacific, Central Pacific, and the Hawaiian Islands and Johnston Atoll.
Countries of Occurrence	American Samoa; Australia; Bahrain; British Indian Ocean Territory; Cambodia; Chile; China; Comoros; Cook Islands; Djibouti; Egypt; Eritrea; Fiji; French Polynesia; Guam; India; Indonesia; Iran, Islamic Republic of; Iraq; Israel; Japan; Jordan; Kenya; Kiribati; Kuwait; Madagascar; Malaysia; Maldives; Marshall Islands; Mauritius; Mayotte; Micronesia, Federated

	States of ; Mozambique; Myanmar; Nauru; New Caledonia; Niue; Northern Mariana Islands; Oman; Pakistan; Palau; Papua New Guinea; Philippines; Pitcairn; Qatar; Réunion; Samoa; Saudi Arabia; Seychelles; Singapore; Solomon Islands; Somalia; South Africa; Sri Lanka; Sudan; Taiwan, Province of China; Tanzania, United Republic of; Thailand; Tokelau; Tonga; Tuvalu; United Arab Emirates; United States Minor Outlying Islands; Vanuatu; Viet Nam; Wallis and Futuna; Yemen
National distribution	<p>Although exact species distribution is unknown, presence records for the species are available in literature and the genus has been observed in every atoll.</p> <p>Data confirmed occurrences of the genus are available from every atoll. Based on known observations, available data, species characteristics it is estimated that there is possibility for the genus and the species to occur on shallow reef systems in the North, Central and South of the Maldives. Depending on depth, the species may not be found.</p>



4. Population:

Summary	Species specific population information is not available. Population estimates and distribution are derived based on best available generic information.
Population Size	<p>Global population is generalized as “common”. National species population size is unknown.</p> <p>A ten-year generic mean, the mean percent cover of the genus over a ten year period of time, was calculated for each genus. This was used as a proxy for the potential species population size. The ten-year generic percent cover mean derived from available data is 0.68 ± 0.04 % (\pm S.E). Recent estimates indicate genus cover is between 0.06 – 1.7 % across most atolls.</p>

Population Trend	Decreasing
Generation length	Estimates for coral generation lengths are based on Carpenter <i>et al.</i> 2008, as follows. The coral colony is considered to be the mature individual, as it typically lives, is injured, or dies as a unit. The average age of natural survival of a coral colony was defined as the average age of a mature individual, or one generation length. Based on available knowledge of coral species' biology and life history, this was determined to be 10 years. Therefore, population reductions are estimated over 30 years, representing three generation lengths.
Continuing Decline in Number of Mature Individuals?	Unknown
Extreme Fluctuations?	No
Severely Fragmented?	No. Population assumed to be cohesive based on generic distribution although degree of connectivity between atolls is unknown.

5. Habitat and Ecology:

Summary	Frequently found between 0 m and 30 m depths, the species occurs in a wide range of shallow reefs habitats, including lagoons, seagrass beds, reef flat as well as turbid waters. Mainly represented by encrusting growth morphologies it can also be found on patchy reefs. Patch reefs are small isolated reef formations that are frequently found within the atoll lagoon.
Systems (terrestrial / freshwater / marine)	Marine
Continuing Decline in Area, Extent, and/or Quality of Habitat?	Yes. There is continuous loss of area, extent and habitat quality stemming from coastal development projects and increase in frequency and magnitude of climate change impacts including extreme temperature fluctuations.
A Migratory Species?	No

6. Use and Trade:

Is the species used or traded?	No
Summary	-

7. Threats:

<p>Summary</p>	<p>The understanding of threats to this species are predominantly based on genus-level threats. The genus is known to be vulnerable to sedimentation, bleaching, predation by corallivores (e.g., <i>Culcita</i>) and disease (WS). The species is affected by overall declines in ecosystem and habitat quality. Therefore, as the species is found in shallow reef habitats (<30 m), it is expected that the species is particularly vulnerable to the impacts of coastal development projects and of climate change. Finally, this is a brooder and therefore has a smaller sexual reproductive output and limited dispersal capacity.</p> <p>The projected date of onset of Annual Severe Bleaching (ASB) is the date after which the capacity of coral reefs to recover from repeated bleaching events is expected to significantly reduce - resulting in fundamental, permanent population changes (UNEP 2017). The calculation of ASB by UNEP (2020) is restricted to a 30 m depth, due to the dataset used for the spatial analysis.</p> <p>We calculated the mean year of ASB onset, assuming 0° coral adaptation (UNEP 2020) to climate change, for areas occupied by the genus. A 0° adaptation (i.e. no adaptation) to climate change was applied as a precautionary approach:</p> <ul style="list-style-type: none"> - as the capacity of coral to adapt to climate change is not yet well understood - as there are other climate change impacts that may compound population decline and affect population refuges, and - to account for impacts from coastal development that could potentially exacerbate the effect of bleaching prompted by climate change. <p>Based on this analysis, <i>L. purpurea</i> is expected to undergo ASB starting in 2041.</p>				
<p>Major Threats to the Species (using IUCN Threat Classification Scheme)</p>	<p>Residential & commercial development:</p> <ol style="list-style-type: none"> 1.1. Housing & urban areas 1.2. Commercial & industrial areas 1.3. Tourism & recreational areas <p>Natural system modifications:</p> <ol style="list-style-type: none"> 7.3. Other ecosystem modifications <p>Invasive & other problematic species, genes & disease:</p> <ol style="list-style-type: none"> 8.2. Problematic native species/diseases <ol style="list-style-type: none"> 8.2.1. Unspecified species/disease <p>Climate change & severe weather:</p> <ol style="list-style-type: none"> 11.1. Habitat shifting & alteration 11.2. Temperature extremes 11.4. Storms & flooding 				
<p>Species vulnerability modifiers</p>	<p>Coral abundance and distribution data is only available at the genus level. Species characteristics were used to evaluate how susceptible each species is to treats. This data was gathered from the IUCN SSC Coral Working Group's Global Traits Database, and available literature. Each species was assigned a vulnerability score for each trait (3 = high vulnerability; 2 = medium vulnerability; 1 = low vulnerability). This species was scored as follows.</p> <table border="1" data-bbox="424 1944 1267 2065"> <thead> <tr> <th>Trait</th> <th>Vulnerability score</th> </tr> </thead> <tbody> <tr> <td>Susceptibility to bleaching</td> <td>3</td> </tr> </tbody> </table>	Trait	Vulnerability score	Susceptibility to bleaching	3
Trait	Vulnerability score				
Susceptibility to bleaching	3				

Resistance to bleaching	3
Recovery from bleaching or disease	3
Resistance to disease	3
Susceptibility to predation	2
Recovery from sedimentation	2
Resistance to sedimentation	2
Susceptibility to sedimentation	2
<i>Average</i>	3

8. Conservation and Research:

Summary	<p>The species is listed as “Near Threatened” on the IUCN Red List of Threatened Species and is included within CITES Appendix II. Collection, killing and export of live and dead scleractinian corals is illegal under Maldivian fisheries law. Distribution of the species falls under multiple national MPAs though not all the MPAs are properly managed. Species distribution and population information would be required to better manage and protect the species and further species-specific research for threats would support development and implementation of necessary legislature</p>
Conservation Actions/Research in Place	Globally, the species is listed within CITES Appendix II and listed as Near Threatened within the IUCN Red List of Threatened Species.
Conservation Actions Needed	<p>Recommended measures for conserving this species include research in taxonomy, abundance and trends, ecology and habitat status, threats and resilience to threats. Restoration action may include artificial reefs to consolidate substrate and enhance the growth of encrusting corals as this species.</p> <ol style="list-style-type: none"> 1. Land/water protection <ol style="list-style-type: none"> 1.1. Site/area protection 1.2. Resource & habitat protection 2. Land/water management <ol style="list-style-type: none"> 2.1. Site/area management 2.3. Habitat & natural process restoration 4. Education & awareness <ol style="list-style-type: none"> 4.1. Formal education 4.2. Training 4.3. Awareness & communications
Research Needed	<ol style="list-style-type: none"> 1. Research <ol style="list-style-type: none"> 1.2. Population size, distribution and past trends 1.3. Life history and ecology

	1.5. Threats 1.6. Conservation actions 3. Monitoring 3.1. Population trends 3.4. Habitat trends
--	---

B. Assessment

Criterion A		
Generation Length	10 years	
3 Generations	30 years	
Reduction in population size?	Yes	
Start Date of Reduction	2022	
End Date of Reduction	2052	
Rate of Reduction (%)	80-90%	
Meets Criteria Thresholds?	A1	
	A2	
	A3	Projected 80-90% reduction over the next three generations (CR A3)
	A4	
Reduction Based on Which Sub-criteria?	a	
	b	
	c	Decline in habitat quality associated with the impacts of climate change
	d	
	e	Mortality due to repeated bleaching

Assessment Under Criterion A	CR A3ce
Reasoning	<p>Our assessment of future population reduction is based on the projected date of onset of ASB. <i>L. purpurea</i> is expected to experience ASB starting in 2041.</p> <p>The species ranges between 0 m and 30 m. Depth and depth range is considered in our assessment, as populations at depths shallower than 30 m are expected to experience greater temperature fluctuations and extreme temperatures (Riegl and Piller 2003), and therefore decline more quickly.</p> <p>Therefore, total population decline is anticipated over 100% of the species' depth range. In addition, the species is considered to have low resistance to bleaching, with high susceptibility to the same threat based on literature review and scientific expertise. Similarly, it is susceptible to diseases and predation.</p> <p>Therefore, this species is projected to undergo a population reduction of 80-90% over the next three generations.</p>

Criterion B		
AOO		-
EOO		-
Meets Criteria Thresholds?	B1	-
	B2	-
Severely Fragmented?		-
No. Locations		-
Threat used to calculate locations		-
Continuing Decline?	(i) EOO	-
	(ii) AOO	-
	(iii) Habitat	-
	(iv) Locations / Subpopulations	-
	(v) Mature Individuals	-
Extreme	(i) EOO	-

Fluctuations?	(ii) AOO	-
	(iii) Locations / Subpopulations	-
	(iv)Mature Individuals	-
Assessment Under Criterion B		DD

Criterion C		
No. Mature Individuals		Unknown
Continuing Decline in Population Size?		-
Is Rate of Decline Known?		-
Generation Length		-
C1	Meets Thresholds for Rate of Decline?	-
	Rate of Decline (%)	-
	Time Period of Decline	-
C2	(a) (i) No. Mature Individuals in Each Subpopulation	-
	(a) (ii) % Mature Individuals in one Sub-population	-
	(b) Extreme Fluctuations in No. Mature Individuals?	-
Assessment Under Criterion C		DD

Criterion D		
No. Mature Individuals		Unknown
Meets Criteria Thresholds?		-
VUD2	AOO	-

	No. Locations	-
	Plausible Future Threat That Would Quickly Drive Taxon to CR or EX	-
	Plausible Future Threat That Would Quickly Drive Taxon to VU or EN	-
Assessment Under Criterion D		DD

Criterion E	
Has a Quantitative Analysis Been Conducted?	No
Type of Quantitative Analysis	-
Generation Length	-
Probability of Extinction within 100 Years	-
Probability of Extinction within 20 Years / 5 Generations (whichever is longer)	-
Probability of Extinction within 10 Years / 3 Generations (whichever is longer)	-
Assessment Under Criterion E	DD

Preliminary Assessment		CR A3ce
Regional Adjustment	Up-list, Down-list, or No Change?	No change
	Justification for Regional Adjustment	No likely interaction with individuals outside the region
Final Assessment		CR A3ce
Narrative Justification for Assessment:		

Leptastrea purpurea is likely distributed throughout all the atolls of the Maldives, with a depth range of 0-30 m. Coral species found at depths of less than 30 m are more exposed to extreme and fluctuating water temperatures associated with the impacts of climate change, which can lead to population reductions.

The most significant known threat to this species is habitat loss and degradation, due to a combination of processes. The species has been identified as being susceptible to bleaching, to predation, and disease.

Although the current population size and past rate of decline are not known, it is anticipated that the population will undergo a severe decline in the future based on the impacts of climate change on its local range. Based on published climate model-based bleaching assessments published by UNEP (2017, 2020), it is estimated that Annual Severe Bleaching (ASB) will commence in the year 2041 over this species' distribution, affecting 100% of its depth range. ASB is expected to constitute a 90-100% reduction in population size.

Therefore, based on the best available data, we project a more than 80% decline in population size within the next three generations (2022-52), prompted by degradation in habitat quality associated with the impacts of climate change. This meets the threshold for CR A3c.

C. References

- Arrigoni, R., Berumen, M. L., Mariappan, K. G., Beck, P. S., Hulver, A. M., Montano, S., ... & Benzoni, F. (2020). Towards a rigorous species delimitation framework for scleractinian corals based on RAD sequencing: the case study of *Leptastrea* from the Indo-Pacific. *Coral Reefs*, 39(4), 1001-1025.
- Carilli, J. E., Bolick, L., Marx, D. E., Smith, S. H., & Fenner, D. (2020). Coral bleaching variability during the 2017 global bleaching event on a remote, uninhabited island in the western Pacific: Farallon de Medinilla, Commonwealth of the Northern Mariana Islands. *Bulletin of Marine Science*, 96(4), 785-802.
- Carpenter, K.E., Abrar, M., Aeby, G., Aronson, R.B., Banks, S., Bruckner, A., Chiriboga, A., Cortés, J., Delbeek, J.C., DeVantier, L., Edgar, G.J., Edwards, A.J., Fenner, D., Guzmán, H.M., Hoeksema, B.W., Hodgson, G., Johan, O., Licuanan, W.Y., Livingstone, S.R., Lovell, E.R., Moore, J.A., Obura, D.O., Ochavillo, D., Polidoro, B.A., Precht, W.F., Quibilan, M.C., Reboton, C., Richards, Z.T., Rogers, A.D., Sanciangco, J., Sheppard, A., Sheppard, C., Smith, J., Stuart, S., Turak, E., Veron, J.E.N., Wallace, C., Weil, E. and Wood, E. 2008. One-Third of Reef-Building Corals Face Elevated Extinction Risk from Climate Change and Local Impacts. *Science*. 25 July 2008: 560-563. DOI: 10.1126/science.1159196. Supporting online material: www.sciencemag.org/cgi/content/full.1159196.
- UNEP 2017. Coral Bleaching Futures - Downscaled projections of bleaching conditions for the world's coral reefs, implications of climate policy and management responses. United Nations Environment Programme, Nairobi, Kenya.
- UNEP 2020. Projections of future coral bleaching conditions using IPCC CMIP6 models: climate policy implications, management applications, and Regional Seas summaries. United Nations Environment Programme, Nairobi, Kenya.
- van Woesik, R., Sakai, K., Ganase, A., & Loya, Y. J. M. E. P. S. (2011). Revisiting the winners and the losers a decade after coral bleaching. *Marine Ecology Progress Series*, 434, 67-76.