



Stylophora pistillata

Hood coral, smooth cauliflower coral

Amir, H.

Citation: Amir, H., 2022. *Stylophora pistillata*. Hard coral, smooth cauliflower coral. The Maldives Red List of Threatened Species.

Maldives National Red List Assessment *Stylophora pistillata*

A. Background Information

1. Assessment Information

Assessor Name(s)	Hana Amir
Date of Assessment	10 February 2022
Contributors	Phillipa Roe, Simone Montano, Ilham Mohamed
Institutional contributors	Ministry of Environment, Climate Change and Technology; Maldives Marine Research Institute; IUCN Project REGENERATE; Maldives Allen Coral Atlas; Noo Raajje; Six Senses Laamu
Facilitators	James Tallant
Reviewers	Janice Chanson, Maldives Marine Research Institute

2. Taxonomic Information

Scientific Name	<i>Stylophora pistillata</i> (Esper, 1792)	
Common Name (English)	Hood coral, Smooth Cauliflower coral	
Common Name (Dhivehi)	No species or generic name	
Taxonomy	Order	Scleractinia
	Sub-order	-
	Family	Pocilloporidae
Taxonomic Notes	Originally <i>Madrepora pistillata</i> (Esper, 1792)	

3. Geographic Range

3.1 Summary of Global Distribution

The species ranges from the Red Sea through the Gulf of Aden, throughout the southwest and northwest Indian Ocean, the Arabian gulf, the central Indian Ocean, Australia, southern Japan, South China Sea, West Pacific Ocean and the Central Pacific Ocean (Hoeksema et al., 2008).


3.2 Countries of Occurrence

American Samoa, Australia, Bahrain, British Indian Ocean Territory, Cambodia, Comoros, Cook Islands, Djibouti, Egypt, Eritrea, Fiji, French Polynesia, Guam, India, Indonesia, Iran, Iraq, Israel, Japan, Jordan, Kenya, Kiribati, Kuwait, Madagascar, Malaysia, Maldives, Marshall Islands, Mauritius, Mayotte, Micronesia, Mozambique, Myanmar, Nauru, New Caledonia, Niue, Norfolk Island, North Mariana Islands, Oman, Palau, Papua New Guinea, Philippines, Qatar, Reunion, Samoa, Saudi Arabia, Seychelles, Singapore, Solomon Islands, Somalia, Sri Lanka, Sudan, Taiwan, Thailand,

Tonga, Tuvalu, United Arab Emirates, United States Minor Outlying Islands, Vanuatu, Vietnam, Wallis and Futuna, Yemen (Hoeksema et al., 2008).

3.3 National distribution

The species' distribution is unknown. Most historical records have the genus restricted to the central-southern and southern atolls. The genus has recently been recorded in Shaviyani, Vaavu, Faafu, Dhaalu, Meemu, Laamu, Huvadhu and Addu atoll. Based on observations, data records and species characteristics, it is estimated that species is likely to be found throughout these atolls. As Shaviyani atoll is part of a bigger geographic atoll ("Bodu Thiladhumathi"), the distribution from the Shaviyani record has been extrapolated for the entire structure. The species is restricted to shallow reef systems and may not be found in the deep atoll lagoons.



4. Population:

4.1 Summary

Species specific population information is not available. Population estimates and distribution are derived based on best available generic information.

4.2 Population Size

Global population is generalized as "common" (Devantier, Turak and Szava-Kovats 2020). National population size is unknown.

A ten-year generic mean, the mean percent cover of the genus over a ten year period of time, was calculated for each genus. This was used as a proxy for the potential species population size.

The ten-year generic mean derived from available data is less than $0.23 \pm 0.08\%$ (\pm S.E). However, this is derived from less than 15 data records.

4.3 Population Trend

Global and national past population trends are not yet known. However, populations have been noted to have been more abundant prior to the 1998 bleaching event with recent sightings considered “rare”. The species has been recorded in the early pioneering expeditions of the Maldives (Pichon and Benzoni 2007) but without population information required to assess a past trend. Future population trend is likely to be decreasing.

4.4 Generation length

Estimates for coral generation lengths are based on Carpenter et al. 2008, as follows. The coral colony is considered to be the mature individual, as it typically lives, is injured, or dies as a unit. The average age of natural survival of a coral colony was defined as the average age of a mature individual, or one generation length. Based on available knowledge of coral species’ biology and life history, this was determined to be 10 years. Therefore, population reductions are estimated over 30 years, representing three generation lengths.

4.5 Continuing Decline in Number of Mature Individuals?

Unknown

4.6 Extreme Fluctuations?

No

4.7 Severely Fragmented?

This species is not considered to be severely fragmented. Degree of fragmentation is unknown. However, populations appear to be restricted to a limited number of atolls within the Maldives.

5. Habitat and Ecology:

5.1 Summary

The species is reef habitat generalist which means it can be found in a variety of reef habitats. It is commonly found between 1 m and 15 m. The species can tolerate both strong and exposed habitats (Veron et al., 2022).

5.2 Systems (terrestrial / freshwater / marine)

Marine

5.3 Continuing Decline in Area, Extent, and/or Quality of Habitat?

Yes. There is continuous loss of area, extent and habitat quality stemming from coastal development projects and increase in frequency and magnitude of climate change impacts including extreme temperature fluctuations.

5.4 A Migratory Species?

No

6. Use and Trade

6.1 Is the species used or traded?

Yes

6.2 Summary

This species is targeted and harvested for the aquarium trade. This harvesting may represent a threat to the species.

7. Threats

7.1 Summary

The understanding of threats to this species are predominantly based on genus-level threats. The species is vulnerable to sedimentation (Ammar et al., 2011), bleaching (Bhagooli and Hidaka 2004; Fitt et al., 2009; Yakovleva and Hidaka 2004), predation, and coral disease (Aeby et al. 2021). Additionally, the species is slow to recover from bleaching and disease once affected.

The projected date of onset of Annual Severe Bleaching (ASB) is the date after which the capacity of coral reefs to recover from repeated bleaching events is expected to significantly reduce - resulting in fundamental, permanent population changes (UNEP 2017). The calculation of ASB by UNEP (2020) is restricted to a 30 m depth, due to the dataset used for the spatial analysis.

We calculated the mean year of ASB onset, assuming 0° coral adaptation (UNEP 2020) to climate change, for areas occupied by the genus. A 0° adaptation (i.e. no adaptation) to climate change was applied as a precautionary approach:

- as the capacity of coral to adapt to climate change is not yet well understood
- as there are other climate change impacts that may compound population decline and affect population refuges, and
- to account for impacts from coastal development that could potentially exacerbate the effect of bleaching prompted by climate change.

Based on this analysis, *S. pistillata* is expected to undergo ASB starting in 2040.

Though not formally recorded within the Maldives, the species is one that is targeted and harvested for the aquarium trade globally.

7.2 Major Threats to the Species (using IUCN Threat Classification Scheme)

1. Residential & commercial development:

1.1. Housing & urban areas

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.1. Ecosystem conversion

1.2. Ecosystem degradation

1.3. Indirect ecosystem effects

2. Species stresses

2.2. Species disturbance

2.3. Indirect species effects

2.3.7. Reduced reproductive success

1.2. Commercial & industrial areas

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.1. Ecosystem conversion

1.2. Ecosystem degradation

1.3. Indirect ecosystem effects

2. Species stresses

2.2. Species disturbance

2.3. Indirect species effects

2.3.7. Reduced reproductive success

1.3. Tourism & recreational areas

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.1. Ecosystem conversion

1.2. Ecosystem degradation

1.3. Indirect ecosystem effects

2. Species stresses

2.2. Species disturbance

2.3. Indirect species effects

2.3.7. Reduced reproductive success

5. Biological resource use:

5.4. Fishing & harvesting aquatic resource

5.4.3. Unintentional effects: subsistence/small scale

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

7. Natural system modifications:

7.3. Other ecosystem modifications

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.1. Ecosystem conversion

- 1.2. Ecosystem degradation
- 1.3. Indirect ecosystem effects

2. Species stresses

- 2.2. Species disturbance
- 2.3. Indirect species effects

- 2.3.7. Reduced reproductive success

8. Invasive & other problematic species, genes & disease:

8.2. Problematic native species/diseases

8.2.1. Unspecified species/disease

2. Species stresses

- 2.1. Species mortality
- 2.2. Species disturbance
- 2.3. Indirect species effects

- 2.3.2. Competition

- 2.3.7. Reduced reproductive success

11. Climate change & severe weather:

11.1. Habitat shifting & alteration

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

- 1.2. Ecosystem degradation

2. Species stresses

- 2.1. Species mortality
- 2.2. Species disturbance
- 2.3. Indirect species effects

- 2.3.2. Competition

- 2.3.7. Reduced reproductive success

11.2. Temperature extremes

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

- 1.2. Ecosystem degradation

2. Species stresses

- 2.1. Species mortality
- 2.2. Species disturbance

2.3. Indirect species effects

2.3.2. Competition

2.3.7. Reduced reproductive success

11.4. Storms & flooding

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

7.3 Species vulnerability modifiers

Coral abundance and distribution data is only available at the genus level. Species characteristics were used to evaluate how susceptible each species is to treats. This data was gathered from the IUCN SSC Coral Working Group's Global Traits Database, and available literature. Each species was assigned a vulnerability score for each trait (3 = high vulnerability; 2 = medium vulnerability; 1 = low vulnerability). This species was scored as follows.

Trait	Vulnerability score
Susceptibility to bleaching	3
Resistance to bleaching	3
Recovery from bleaching or disease	3
Resistance to disease	3
Susceptibility to predation	2
Recovery from sedimentation	3
Resistance to sedimentation	3
Susceptibility to sedimentation	3
<i>Average</i>	3

8. Conservation and Research

8.1 Summary

The species is listed as "Near Threatened" on the IUCN Red List of Threatened species and is listed on CITES Appendix II. Collection, killing and export of live and dead scleractinian corals is illegal under Maldivian fisheries law. Distribution of the species falls under multiple national MPAs though not all the MPAs are properly managed. Species distribution and population information would be required to better manage and protect the species and support the development and

implementation of necessary legislatures. Targeted studies are required to understand whether and to what degree the species is locally exploited for the aquarium trade or propagations projects.

8.2 Conservation Actions/Research in Place

Globally, as a coral the species is listed within CITES Appendix II. It is listed as “Near Threatened” within the IUCN Red List of Threatened Species.

Nationally, collection, killing, and export of both live and dead scleractinian corals (under phylum Cnidaria) are illegal in the Maldives under the Maldives Fisheries Law 2020/R-75. The distribution of the genus, and likely the species, overlaps multiple MPAs including Baa atoll biosphere reserve and SAMPA.

8.3 Summary

Despite laws and regulations protecting coral species within the country, implementation of such laws and regulations are a challenge. There is a need to create infrastructure and capacity that would enable relevant laws and regulations to be enforced whilst educating the general public that they exist. Educational and training needs extend to more educational opportunities in the marine field as well as skill training opportunities that can support research, further education, institutional development, awareness and communication efforts. Moreover, with the spatial distribution of maldivian reefs and the corals that are found within them, there is an urgent need to foster intra-governmental as well as inter-agency and inter-organizational cooperation and partnerships to mediate limitations related to capacity and resources. Areas of high abundance of specific species along with areas of high coral cover and resilience need to be identified and managed if not fully protected to ensure sustainable use and longevity. Management includes mitigating impacts from coastal development and resolving conflict of multi-use resources. Further, with increasing interest in restoration of reefs, there is a need to properly manage species harvest for projects to ensure that wild stocks are not decimated and species are able to recover.

1. Land/water protection

1.1. Site/area protection

1.2. Resource & habitat protection

2. Land/water management

2.1. Site/area management

2.3. Habitat & natural process restoration

3. Species management

3.1. Species management

3.1.1. Harvest management

3.2. Species recovery

3.4. *Ex situ* conservation

3.4.1. Captive breeding/artificial propagation

3.4.2. Genome resource bank

4. Education & awareness

4.1. Formal education

4.2. Training

4.3. Awareness & communications

5. Law & policy

5.3. Private sector standards & codes

5.4. Compliance & enforcement

5.4.1. Scale unspecified

7. External capacity building

7.1. Institutional & civil society development

7.2. Alliance & partnership development

7.3. Conservation finance

8.4 Conservation Actions Needed

1. Research

1.2. Population size, distribution and past trends

1.3. Life history and ecology

1.5. Threats

1.6. Conservation actions

3. Monitoring

3.1. Population trends

3.4. Habitat trends

B. Assessment

9. CRITERION A

Criterion A

Generation Length	10 years	
3 Generations	30 years	
Reduction in population size?	Yes	
Start Date of Reduction	2022	
End Date of Reduction	2052	
Rate of Reduction (%)	80-90%	
Meets Criteria Thresholds?	A1	-
	A2	-
	A3	Projected 80-90% reduction over the next three generations (CR A3)
	A4	-
Reduction Based on Which Sub-criteria?	a	-
	b	-
	c	Decline in habitat quality associated with climate change.
	d	-
	e	Mortality due to repeated bleaching

Assessment Under Criterion A

CR A3ce

Reasoning Our assessment of future population reduction is based on the projected date of onset of ASB. *S. pistillata* is expected to experience ASB starting in 2040. The species has a depth range of 1-15 m and is fully restricted to depths shallower than 30 m. Depth and depth range is considered in our assessment,

as populations at depths shallower than 30m are expected to experience greater temperature fluctuations and extreme temperatures (Riegl and Piller 2003), and therefore decline more quickly.

The species range is also limited with records from only a few atolls.

The species is considered to be relatively vulnerable to bleaching (Bhagooli and Hidaka 2004; Fitt et al., 2009; Yakovleva and Hidaka 2004), sedimentation (Ammar et al., 2011) and coral disease (Aeby et al., 2021) with high susceptibility and low resilience based on literature review and scientific expertise. Therefore, total population decline is anticipated over 100% of the species' depth range.

Therefore, this species is projected to undergo a population reduction of 80-90% over the next three generations (2022-2052).

10. CRITERION B

Criterion B

AOO		-
EOO		-
Meets Criteria Thresholds?	B1	-
	B2	-
Severely Fragmented?		-
No. Locations		-
Threat used to calculate locations		-
Continuing Decline?	(i) EOO	-
	(ii) AOO	-
	(iii) Habitat	-
	(iv) Locations / Subpopulations	-
	(v) Mature Individuals	-

Extreme Fluctuations?	(i) EOO	-
	(ii) AOO	-
	(iii) Locations / Subpopulations	-
	(iv) Mature Individuals	-

Assessment Under Criterion B DD

11. CRITERION C

Criterion C

No. Mature Individuals		Unknown
Continuing Decline in Population Size?		-
Is Rate of Decline Known?		-
Generation Length		-
C1	Meets Thresholds for Rate of Decline?	-
	Rate of Decline (%)	-
	Time Period of Decline	-
C2	(a) (i) No. Mature Individuals in Each Subpopulation	-
	(a) (ii) % Mature Individuals in one Sub-population	-
	(b) Extreme Fluctuations in No. Mature Individuals?	-

Assessment Under Criterion C DD

12. CRITERION D

Criterion D

No. Mature Individuals	Unknown
Meets Criteria Thresholds?	-
VUD2 AOO	-
No. Locations	-
Plausible Future Threat That Would Quickly Drive Taxon to CR or EX	-
Plausible Future Threat That Would Quickly Drive Taxon to VU or EN	-

Assessment Under Criterion D DD

13. CRITERION E

Criterion E

Has a Quantitative Analysis Been Conducted?	No
Type of Quantitative Analysis	-
Generation Length	--
Probability of Extinction within 100 Years	-
Probability of Extinction within 20 Years / 5 Generations (whichever is longer)	-

Probability of Extinction within 10 Years / 3 Generations (whichever is longer) -

Assessment Under Criterion E DD

14. Preliminary and Final Assessment

Preliminary Assessment CR A3ce

Regional Adjustment	Up-list, Down-list, or No Change?	No change
	Justification for Regional Adjustment	No likely interaction with individuals outside the region

Final Assessment CR A3ce

Narrative Justification for Assessment:

Stylophora pistillata appears to be distributed in a few atolls of the Maldives, with a depth range of 1-15 m. Coral species found at depths of less than 30 m are more exposed to extreme and fluctuating water temperatures associated with the impacts of climate change, which can lead to population reductions.

This species is believed to be impacted by threats of sedimentation, bleaching, predation by corallivores, and disease. It has been identified as being vulnerable to bleaching (with high susceptibility and low resistance). In addition, it has been seen to be slow to recover from bleaching and disease once affected.

Although its population size is not known, we anticipate that this species will undergo a severe decline in the future based on the impacts of climate change on its local range. Based on published climate model-based bleaching assessments published by UNEP (2017, 2020), Annual Severe Bleaching (ASB) is likely to commence in 2040 over this species' distribution, affecting 100% of its depth range. ASB reduces corals' ability to recover from repeated bleaching events, resulting in fundamental, permanent population changes (UNEP 2017). ASB is expected to constitute at least a 80-90% reduction in population size. Therefore, we project a 80-90% population reduction over the next three generations (2022-2052), resulting from degradation in habitat quality associated with the impacts of climate change. This meets the threshold for CR A3c.

C. References

- Aeby, G. S., Shore, A., Jensen, T., Ziegler, M., Work, T., & Voolstra, C. R. (2021). A comparative baseline of coral disease across the central Red Sea. *bioRxiv*.
- Ammar, M. S., Obuid-Allah, A. H., & Al-Hammady, M. A. (2011). Corals differential susceptibilities to bleaching along the Red Sea Coast, Egypt. *Nusantara Bioscience*, 3(2).
- Bhagooli, R., & Hidaka, M. (2004). Photoinhibition, bleaching susceptibility and mortality in two scleractinian corals, *Platygyra ryukyuensis* and *Stylophora pistillata*, in response to thermal and light stresses. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 137(3), 547-555.
- Carpenter, K.E., Abrar, M., Aeby, G., Aronson, R.B., Banks, S., Bruckner, A., Chiriboga, A., Cortés, J., Delbeek, J.C., DeVantier, L., Edgar, G.J., Edwards, A.J., Fenner, D., Guzmán, H.M., Hoeksema, B.W., Hodgson, G., Johan, O., Licuanan, W.Y., Livingstone, S.R., Lovell, E.R., Moore, J.A., Obura, D.O., Ochavillo, D., Polidoro, B.A., Precht, W.F., Quibilan, M.C., Reboton, C., Richards, Z.T., Rogers, A.D., Sanciangco, J., Sheppard, A., Sheppard, C., Smith, J., Stuart, S., Turak, E., Veron, J.E.N., Wallace, C., Weil, E. and Wood, E. 2008. One-Third of Reef-Building Corals Face Elevated Extinction Risk from Climate Change and Local Impacts. *Science*. 25 July 2008: 560-563. DOI: 10.1126/science.1159196. Supporting online material: www.sciencemag.org/cgi/content/full.1159196.
- DeVantier, L., & Turak, E. (2017). Species richness and relative abundance of reef-building corals in the Indo-West Pacific. *Diversity*, 9(3), 25.
- DeVantier, L., Turak, E., & Szava-Kovats, R. (2020). Species Richness and Abundance of Reef-Building Corals in the Indo-West Pacific: The Local–Regional Relation Revisited. *Frontiers in Marine Science*, 7, 487.
- Fitt, W. K., Gates, R. D., Hoegh-Guldberg, O., Bythell, J. C., Jatkar, A., Grottoli, A. G., ... & Lesser, M. P. (2009). Response of two species of Indo-Pacific corals, *Porites cylindrica* and *Stylophora pistillata*, to short-term thermal stress: the host does matter in determining the tolerance of corals to bleaching. *Journal of experimental marine biology and ecology*, 373(2).
- Hoeksema, B. W.; Cairns, S. (2022). World List of Scleractinia. *Stylophora pistillata* (Esper, 1792). Accessed through: World Register of Marine Species at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=206982> on 2022-02-05
- Hoeksema, B.W., Rogers, A. & Quibilan, M.C. 2014. *Stylophora pistillata*. *The IUCN Red List of Threatened Species* 2014: e.T133685A54306217. <https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T133685A54306217.en>. Accessed on 04 March 2022.
- Pichon, M., & Benzoni, F. (2007). Taxonomic re-appraisal of zooxanthellate Scleractinian Corals in the Maldivian Archipelago. *Zootaxa*, 33(1441), 21–33. <https://doi.org/10.11646/zootaxa.1441.1.2>
- Rasheed, S., Warder, S. C., Plancherel, Y., & Piggott, M. D. (2021). An Improved Gridded Bathymetric Data Set and Tidal Model for the Maldives Archipelago. *Earth and Space Science*. <https://doi.org/10.1029/2020EA001207>
- Riegl, B. and Piller, W.E. 2003. Possible refugia for reefs in times of environmental stress. *International Journal of Earth Sciences* 92(4):520-531.
- UNEP 2017. Coral Bleaching Futures - Downscaled projections of bleaching conditions for the world's coral reefs, implications of climate policy and management responses. United Nations Environment Programme, Nairobi, Kenya
- UNEP 2020. Projections of future coral bleaching conditions using IPCC CMIP6 models: climate policy implications, management applications, and Regional Seas summaries. United Nations Environment

Programme, Nairobi, Kenya

Veron J.E.N., Stafford-Smith M.G., Turak E. and DeVantier L.M. (2022). Corals of the World. Accessed 9 February 2022, Version 0.01 (Beta). [http://coralsoftheworld.org/v0.01\(Beta\)](http://coralsoftheworld.org/v0.01(Beta)). (To go to the current version access: <http://coralsoftheworld.org>)

Yakoleva, I., & Hidaka, M. (2004). Different effects of high temperature acclimation on bleaching-susceptible and tolerant corals. *Symbiosis*.